



2170 West State Road 434, Suite 190
 Longwood, FL 32779
 Phone: 407-990-1921 | Fax: 855-537-4411
 Website: <https://heartveinvascular.com>

Superficial Venous Reflux Treatment

Superficial venous reflux is a condition that develops when the valves that usually keep blood flowing out of your legs become damaged or diseased. This causes blood to pool in your legs. Common symptoms of superficial venous reflux include pain, swelling, leg heaviness and fatigue, as well as varicose veins in your legs. The Closure procedure is a minimally invasive treatment for superficial venous reflux. A thin catheter is inserted into the vein through a small opening. The catheter delivers radiofrequency (RF) energy to the vein wall, causing it to heat, collapse, and seal shut. Since valves can't be repaired, the only alternative is to re-route blood flow through healthy veins. Traditionally, this has been done by surgically removing (stripping) the troublesome vein from your leg. The Closure procedure provides a less invasive alternative to vein stripping by simply closing the problem vein instead. Once the diseased vein is closed, other healthy veins take over and empty blood from your legs. In the Closure procedure, there is no need for groin surgery. Instead, the vein remains in place and is closed using a special (Closure) catheter inserted through a small puncture. This may eliminate the bruising and pain often associated with vein stripping (i.e., that may result from the tearing of side branch veins while the saphenous vein is pulled out). Vein stripping is usually performed in an operating room, under a general anesthetic, while the Closure procedure is performed on an outpatient basis, typically using local or regional anesthesia. The Closure procedure takes approximately 45-60 minutes, though patients normally spend 2-3 hours at the medical facility due to normal pre- and post-treatment procedures. Many patients can resume normal activities immediately.² For a few weeks following the treatment, your doctor may recommend a regular walking regimen and suggest you refrain from very strenuous activities (heavy lifting, for example) or prolonged periods of standing. Most patients report a noticeable improvement in their symptoms within 1-2 weeks following the procedure. Patients report minimal to no scarring, bruising, or swelling following the Closure procedure. As with any medical intervention, potential risks and complications exist with the Closure procedure. All patients should consult their doctors to determine if their conditions present any special risks. Your physician will review potential complications of the Closure procedure at the consultation, and can be reviewed in the safety summary. Potential complications can include: vessel perforation, thrombosis, pulmonary embolism, phlebitis, hematoma, infection, paresthesia (numbness or tingling), and/or skin burn. The vein simply becomes fibrous tissue after treatment. Over time, the vein will gradually incorporate into surrounding tissue. One study reported that 89% of treated veins are indistinguishable from other body tissue one year after the Closure procedure was performed.

Procedure Date: _____ Time: _____

_____ I understand that I will be charged \$130 if I do not cancel or reschedule my procedure at least 2 business days in advance.

_____ I understand that I will not hold Orlando Heart Specialists or Center for Vein and Vascular Disease liable if I do not undergo my procedure as scheduled.

Patient Signature: _____ Today's Date : _____

Patient Name: _____